

# MCQ Geriatric Workshop

Dr. Akhil Kapoor

Associate Professor, Medical Oncology

Tata Memorial Hospital (HBCH &MPMMCC), Varanasi



Q. 1. Which of the following is **NOT** correct about G8 Screening tool?

1. Score lies between 0-17
2. Higher Score: Better health status
3. Score more than 15 should undergo CGA
4. Includes age, food intake, weight loss, mobility, BMI, neuropsychological problems

## Q. 2. Which of the following is FALSE about CARG tool?

1. It is based on data of 500 older pts receiving chemo (Hurria)
2. Poly Vs. Monochemo included as part of risk assessment.
3. Type of cancer is a part of the tool.
4. Administration time is around 20 mins.

## Q. 3. Which is **TRUE** about CARG Vs. CRASH?

1. CARG tool is validated for patients aged  $\geq 70$ .
2. CRASH tool's administration time is shorter.
3. CRASH tool is validated for patients aged  $\geq 65$ .
4. CRASH includes IADL, MMSE, MNA.

## Q. 4. What is NOT correct about the GO2 trial?

1. It was a Phase III trial in older patients with Gastroesophageal cancer patients.
2. Three dose levels of CAPOX with 100, 80 and 60% doses were used.
3. Both PFS and OS were found to be similar in all three dose levels.
4. Reduced doses did not lead to any differences in the QOL.

## Q. 5. Which of the following is CORRECT about GAP70+ Study?

1. Geriatric assessment led to improved survival in the patients with age of 70+ years.
2. Grade 1-2 toxicities were significantly reduced by following Geriatric assessment.
3. Grade 3-5 toxicities were significantly reduced by following Geriatric assessment.
4. Significantly higher number of patients in Geriatric assessment group were offered more aggressive chemotherapy.

# Answers

- 1. 3
- 2. 4
- 3. 4
- 4. 4
- 5. 3